Gauging staggered fermion shift symmetries and lattice anomalies

Simon Catterall^{1 \star} and Arnab Pradhan^{2 \dagger}

Department of Physics, Syracuse University, Syracuse, NY 13244, USA *smcatter@syr.edu *arpradha@syr.edu

May 7, 2024

Abstract

Staggered fermion shift symmetries correspond to translations within the unit cell of a hypercubic lattice. They satisfy an algebra and in four Euclidean dimensions can be related to a discrete subgroup of SU(4) which plays a crucial role in showing that staggered fermions lead to a theory of four degenerate Dirac fermions in the continuum limit. They are associated with the appearance of certain Z_2 valued global parameters. We propose a strategy to try to partially gauge these translation symmetries by allowing these parameters to vary locally in the lattice. To maintain invariance of the action requires the addition of Z_2 valued higher form gauge fields. We show how there is a geometrical obstruction to this procedure in the case of full staggered fermions which is consistent with the existence of a 't Hooft anomaly in the continuum SU(4) flavor symmetry. However no such obstruction is encountered for reduced staggered fermions for which the continuum non-anomalous flavor symmetry is SO(4).

Contents

1	Introduction	2
2	Global shift symmetries of staggered fermions	3
3	Continuum interpretation of the shifts	4
4	Gauging shifts and higher form gauge fields	6
	4.1 Single shifts	6
	4.2 Higher shifts	7
	4.3 Obstructions to gauging - lattice 't Hooft anomalies	8
5	Gauging shifts for reduced staggered fermions	9
6	Summary	10
A	Identities	11
References		12

1 Introduction

In this paper we will be concerned with the properties of staggered lattice fermions. Staggered fermions are usually regarded as discretizations of Dirac fermions but in fact it is more natural to regard them as discrete Kähler-Dirac fermions [1–5]. The Kähler-Dirac equation affords an alternative to the Dirac equation and is written in terms of the exterior derivative and its adjoint [6]. It naturally acts on (Grassmann valued) antisymmetric tensor fields (p-forms) which replace the spinors of the Dirac theory. In four dimensional flat space one can show that a single Kähler-Dirac field gives rise to four degenerate Dirac fields but this correspondence is no longer true in a curved background - for example the Kähler-Dirac operator has exact zero modes on the sphere unlike its Dirac cousin.

Kähler-Dirac fermions are particularly interesting as they can be discretized without breaking a certain global U(1) onsite symmetry and without inducing fermion doubling - there is no analog of the Nielsen-Ninomiya theorem in this case [7]. Furthermore, contrary to the usual folklore that lattice symmetries are necessarily free of anomalies, the lattice Kähler-Dirac theory suffers from a mixed U(1)-gravitational anomaly that breaks U(1) to Z_4 on spaces with the topology of a sphere [8–10]. Attempts to gauge this residual Z_4 symmetry reveal the presence of a 't Hooft anomaly. Cancellation of this 't Hooft anomaly places constraints on the number of fermions needed to obtain massive symmetric phases in theories of Kähler-Dirac or staggered fermions in a variety of dimension [9–14]. In fact if we decompose the Kähler-Dirac field into a set of spinors we find that the anomaly free theories contain precisely the same number of Majorana fermions that are needed to gap boundary states in topological superconductors in any dimension [15–19].

The generator of the U(1) symmetry can also be used to define a projector which acts on a Kähler-Dirac field to remove half the degrees of freedom. In the continuum flat space theory it is straightforward to see that this so-called *reduced* Kähler-Dirac field is in fact a chiral theory comprising a doublet of left and right handed Weyl fields transforming under two independent SU(2) global flavor symmetries [5, 10]. In the case of staggered fermions this projection corresponds to employing reduced staggered fermions. It is thus natural to investigate such theories as a possible way to construct certain classes of chiral lattice gauge theory [10, 20].

If such a strategy is to succeed it is crucial to understand how the existence of certain exact lattice symmetries control the approach to the continuum limit. In particular the massless staggered fermion action is invariant under a group of discrete shift symmetries that correspond to a subgroup of the flavor symmetries of the continuum Dirac theory [21–23]. These shifts correspond to lattice translation within the unit hypercube with appropriately chosen site-dependent phases that leave the action invariant.

To guarantee a chiral continuum theory requires that no bilinear terms coupling staggered fields within the unit hypercube arise since they would correspond to mass terms. Shift invariance is already sufficient to guarantee that no such terms can arise in the effective action but it is also important to understand whether an effective coupling can arise either via an anomaly or through spontaneous symmetry breaking. This requires understanding the non-perturbative structure of the theory and one of the few tools we have at our disposal to probe such issues is an enumeration of all possible 't Hooft anomalies. Thus, the question that naturally arises is whether there is any obstruction to gauging these lattice translation symmetries. This is non-trivial since these symmetries do not act onsite and hence the action cannot be trivially gauged by inserting gauge links. ¹

In this paper we attempt to provide at least a partial gauging of these shift symmetries. We

¹Invariance of the fermion measure under local shifts requires the theory contain multiples of two such reduced fields but this is already guaranteed by cancelling off the Z_4 't Hooft anomaly mentioned earlier.

find a need to invoke higher form gauge fields to do this and further find that the procedure we use will necessarily fail for the full set of shift symmetries of a staggered fermion. However, a reduced staggered fermion does mot suffer from the same obstruction and we tentatively conclude that there are no such 't Hooft anomalies arising from the lattice shift symmetries in this case.

We start with a review of staggered fermions and shift symmetries emphasising the importance of the latter for restoration of continuum flavor symmetry in the continuum limit. We then describe our strategy for gauging these symmetries and the appearance of higher form fields. We articulate the geometrical obstructions to this lattice gauging procedure and comment on the connection to continuum 't Hooft anomalies. Finally we show how a consistent gauging can be obtained for so-called *reduced* staggered fermions. In the final section we describe the consequences of this work to a program of using (reduced) staggered fermions to target a class of chiral gauge theory.

2 Global shift symmetries of staggered fermions

The massless four dimensional staggered fermion action takes the form

$$S = \sum_{x,\mu} \eta_{\mu}(x)\overline{\chi}(x)[\chi(x+\mu) - \chi(x-\mu)]$$
(1)

where $\eta_{\mu}(x) = (-1)^{\sum_{i=1}^{\mu-1} x_i}$ are the usual staggered fermion phases. Consider performing a lattice translation or shift transformation in the direction λ :

$$\chi(x) \to i\alpha_{\lambda}\xi_{\lambda}(x)\chi(x+\lambda)$$

$$\overline{\chi}(x) \to -i\alpha_{\lambda}\xi_{\lambda}(x)\overline{\chi}(x+\lambda)$$
(2)

with $\xi_{\mu}(x) = (-1)^{\sum_{i=\mu+1}^{4} x_i}$ and the global α_{λ} parameter taking its values in the group $Z_2 \in \{1, -1\}$. The action transforms to

$$S' = \sum_{x,\mu} \eta_{\mu}(x) \overline{\chi}(x+\lambda) \xi_{\lambda}(x) [\xi_{\lambda}(x+\mu)\chi(x+\mu+\lambda) - \xi_{\lambda}(x-\mu)\chi(x-\mu+\lambda)]$$
$$= \sum_{x,\mu} \eta_{\mu}(x) \overline{\chi}(x) [\chi(x+\mu) - \chi(x-\mu)] [\eta_{\mu}(\lambda)\xi_{\lambda}(\mu)]$$
(3)

But it is easy to see from the definitions of the phase factors that $\eta_{\mu}(\lambda)\xi_{\lambda}(\mu) = 1$ (see the appendix) and so the action is invariant under this translation [21–23]. Notice that two successive transformations in the same direction accomplish a simple translation $\chi(x) \rightarrow \chi(x+2\lambda)$ (up to a trivial change of sign of χ).

You can compound these shifts to build other discrete symmetries. For example the following double shift transformation in the ρ , λ directions (with $\rho \neq \lambda$) is also a symmetry

$$\chi(x) \to \alpha_{\lambda\rho} \xi_{\lambda}(x) \xi_{\rho}(x+\lambda) \chi(x+\lambda+\rho)$$

$$\overline{\chi}(x) \to \alpha_{\lambda\rho} \xi_{\lambda}(x) \xi_{\rho}(x+\lambda) \overline{\chi}(x+\lambda+\rho) \quad \text{no sum on } \rho \text{ and } \lambda \text{ and } \alpha_{\lambda\rho} \in \mathbb{Z}_{2}$$
(4)

Using the relation (see the appendix)

$$\xi_{\mu}(x)\xi_{\nu}(x+\mu) + \xi_{\nu}(x)\xi_{\mu}(x+\nu) = 2\delta_{\mu\nu}$$
(5)

one can show that $\alpha_{\lambda\rho} = -\alpha_{\rho\lambda}$ showing that the $\alpha_{\rho\lambda}$ behave like Z_2 valued constant antisymmetric tensors. These double shifts also obey an algebra. For example consider the combination of two such double shifts characterized by (μ, ν) and (ν, λ) (we set the α parameters to

unity for simplicity)

$$\chi(x) \to \xi_{\mu}(x)\xi_{\nu}(x+\mu)\xi_{\nu}(x+\mu+\nu)\xi_{\lambda}(x+\mu+2\nu)\chi(x+\mu+2\nu+\lambda)$$

$$\chi(x) \to \xi_{\mu}(x)\xi_{\lambda}(x+\mu)\chi(x+\mu+\lambda+2\nu)$$
(6)

So two such double shifts give rise to a third distinct (μ, λ) double shift plus an ordinary translation by 2ν . The relations given in eqn. 5 and eqn. 6 are reminiscent of the properties of (Euclidean) Dirac matrices and suggest a connection to continuum symmetries. This will turn out to be true and is examined in the next section.

3 Continuum interpretation of the shifts

To get a better understanding of the origin of these shift symmetries it is helpful to assemble all the staggered fields in a hypercube into a matrix fermion

$$\Psi(x) = \frac{1}{8} \sum_{b} \chi(x+b) \gamma^{x+b}$$
(7)

In this expression b_i is a 4-component vector with components $b_i = 0, 1$ and the sum extends over all the points in the unit hypercube with corner x. The notation γ^x denotes $\gamma^x = \gamma_1^{x_1} \gamma_2^{x_2} \gamma_3^{x_3} \gamma_4^{x_4}$. Clearly the matrix field $\Psi(x)$ carries sixteen times as many degrees of freedom as the original staggered field. The natural way to restore the correct counting of degrees of freedom is to consider $\Psi(x)$ as defined on a lattice with twice the lattice spacing – the so-called spin-taste basis of staggered fermions [24]. The various staggered fields within the hypercube can then be interpreted as comprising the spin and flavor components of a series of four Dirac fermions corresponding to the columns of Ψ . This representation can be related to the Kähler-Dirac interpretation of staggered fermions [1,3].

It is possible to write down a continuum action for this matrix fermion that can be naturally discretized to give the staggered fermion action

$$S = \int d^4 x \operatorname{Tr} \left[\overline{\Psi} \gamma^{\mu} \partial_{\mu} \Psi \right]$$
(8)

To see this in detail one merely has to substitute eqn. 7 into the matrix action and replace the derivative by a symmetric difference operator – after doing the traces one obtains the staggered fermion action in eqn. 1. The phase $\eta_{\mu}(x)$ just reflects the net effect of commuting the extra factor of γ^{μ} from the displaced fields $\chi(x \pm \mu)$ through a string of gamma matrices until it multiplies the explicit factor of γ^{μ} in the kinetic operator.

The symmetries of the continuum action in eqn. 8 are now manifest and correspond to the transformations

$$\begin{split} \Psi &\to L \Psi F^{\dagger} \\ \overline{\Psi} &\to F \overline{\Psi} L^{\dagger} \end{split} \tag{9}$$

where *L* are (Euclidean) *SO*(4) Lorentz transformations acting on the spinors that form the columns of Ψ and *F* are *SU*(4) flavor transformations. For the flavor group we can adopt a basis corresponding to the Clifford algebra associated to products of Euclidean Dirac matrices. Thus $F = e^{i \sum_{A} \beta_{A} \Gamma^{A}}$ where $\Gamma^{A} = {\hat{\gamma}_{\mu}, i \hat{\gamma}_{\mu\nu}, i \hat{\gamma}_{5} \hat{\gamma}_{\mu}, \hat{\gamma}_{5}}$.

In general both Lorentz and flavor symmetries are broken once we employ a staggered fermion discretization. While the breaking of Lorentz symmetry to the group of hypercubic rotations is obvious the breaking of flavor symmetries relies on the observation that staggered fermions, like their Kähler-Dirac cousins, are only invariant under a diagonal subgroup of the continuum Lorentz and flavor symmetries corresponding to L = F in eqn. 9. In particular the only flavor symmetries that survive discretization correspond to transformations arising from rotations of the form $e^{\alpha_A i \frac{\pi}{2} \Gamma^A} = i \alpha_A \Gamma^A$ where $\alpha_A \in Z_2$ with the further restriction that the flavor basis matrices $\hat{\gamma}_{\mu}$ are to be identified with the Dirac basis γ_{μ} . For example, taking $\Gamma^A \equiv \gamma^{\lambda}$ the lattice matrix fermion transforms as

$$\Psi(x) \to \Psi(x+\lambda)i\alpha_{\lambda}\gamma^{\lambda}$$

$$\overline{\Psi}(x) \to -i\alpha_{\lambda}\gamma^{\lambda}\overline{\Psi}(x+\lambda)$$
(10)

Using this result it is not hard to see that it is this discrete flavor transformation that gives rise to the staggered shift symmetry described earlier. Using eqn. 7 one finds

$$\Psi' = \frac{1}{8} \sum_{b} \chi(x+b+\lambda) \gamma^{x+b+\lambda} i \alpha_{\lambda} \gamma^{\lambda}$$
$$= \frac{1}{8} \sum_{b} i \alpha_{\lambda} \xi_{\lambda}(x+b) \chi(x+b+\lambda) \gamma^{x+b+\lambda}$$
(11)

where the phase $\xi_{\lambda}(x)$ arises because of commuting γ^{λ} from the right through to its appropriate spot in the string of γ matrices. Clearly, the net effect is to shift $\chi(x) \rightarrow i \alpha_{\lambda} \xi_{\lambda}(x) \chi(x + \lambda)$ - a shift transformation on the staggered fermion. Similarly the double shifts discussed earlier correspond to continuum transformations of the form $e^{\frac{\pi}{2}\alpha_{\lambda\rho}\gamma^{\lambda}\gamma^{\rho}} = \alpha_{\lambda\rho}\gamma^{\lambda}\gamma^{\rho}$.

In four dimensions there are 32 discrete shift symmetries of the staggered action that that can be built using these transformations and their inverses. These correspond to the discrete subgroup Γ_4 of SU(4) given by

$$\Gamma_4 = \{\pm I, \pm \gamma^{\lambda}, \pm i\gamma^{\lambda}\gamma^{\rho}, \pm i\gamma^5\gamma^{\lambda}, \pm \gamma^5\}$$
(12)

It is the existence of this subgroup of lattice shift symmetries for staggered fermions that guarantees restoration of full SU(4) flavor symmetry in the continuum limit – one cannot write down any relevant counterterms which are invariant under this discrete subgroup but which break SU(4) [22].

Actually, there are two independent sets of shift symmetries. Our discussion so far has focused on the transformation

$$\chi(x) \xrightarrow{\Gamma_{4}^{V}} +i\alpha^{\lambda}\xi_{\lambda}(x)\chi(x+\lambda)$$

$$\overline{\chi}(x) \to -i\alpha^{\lambda}\xi_{\lambda}(x)\overline{\chi}(x+\lambda)$$
(13)

corresponding to the continuum transformation of $\Psi \to e^{i\frac{\pi}{2}\alpha_{\lambda}\gamma^{\lambda}}\Psi$ and $\overline{\Psi} \to e^{-i\frac{\pi}{2}\alpha_{\lambda}\gamma^{\lambda}}\overline{\Psi}$. The other set of transformations are given by

$$\chi(x) \xrightarrow{\Gamma_4^{\Lambda}} +i\epsilon(x)\alpha_{\lambda}\xi_{\lambda}(x)\chi(x+\lambda)$$

$$\overline{\chi}(x) \to +i\epsilon(x)\alpha_{\lambda}\xi_{\lambda}(x)\overline{\chi}(x+\lambda)$$
(14)

corresponding to $\Psi \to e^{i\alpha_{\lambda}\frac{\pi}{2}\gamma^{5}\otimes\gamma^{5}\gamma^{\lambda}}\Psi$ and $\overline{\Psi} \to e^{i\alpha_{\lambda}\frac{\pi}{2}\gamma^{5}\otimes\gamma^{5}\gamma^{\lambda}}\overline{\Psi}$. These are the analog of "axial" transformations - where γ^{5} is replaced by $\gamma^{5} \otimes \gamma^{5}$ which has both a right and left action on the matrix fermion. It becomes the site parity operator $\epsilon(x)$ of staggered fermions. Thus the full lattice shift symmetry is $\Gamma_{4}^{V} \times \Gamma_{4}^{A}$.

4 Gauging shifts and higher form gauge fields

4.1 Single shifts

To gauge these symmetries we will let the parameters α depend on position. To start we focus on the single shift symmetry and allow $\alpha_{\lambda} = \alpha_{\lambda}(x)$ (we simplify the expressions by leaving out the factors of i). The transformation is now

$$\chi(x) \to \alpha_{\lambda}(x)\xi_{\lambda}(x)\chi(x+\lambda) \tag{15}$$

What happens when a shift of two lattice spacings is made in the same direction ? In this case

$$\chi(x) \to \alpha_{\lambda}(x)\alpha_{\lambda}(x+\lambda)\chi(x+2\lambda) \tag{16}$$

Since one wants such a shift to be a simple translation this seems to require $\alpha_{\lambda}(x)\alpha_{\lambda}(x+\lambda) = 1$. However, this is a strong constraint since it determines all $\alpha_{\lambda}(x)$ in direction λ from their value on a single slice. However, there is another possibility – namely that the factor $\alpha_{\lambda}(x)\alpha_{\lambda}(x+\lambda)$ can be interpreted as a local Z_2 gauge transformation acting on $\chi(x)$. In this way the two lattice spacing shift would yield a simple translation up to an unobservable Z_2 phase. Of course, this requires the introduction of a Z_2 link gauge field $L_{\mu}(x)$ to keep the action invariant. The action is then given by

$$S = \sum_{x,\mu} \eta_{\mu}(x)\overline{\chi}(x) \left[L_{\mu}(x)\chi(x+\mu) - L_{\mu}(x-\mu)\chi(x-\mu) \right]$$
(17)

A Wilson plaquette term can be added for the link field $L_{\mu}(x)$

$$S_L = -\beta_L \sum_{x,\mu < \nu} F_{\mu\nu} \tag{18}$$

where

$$F_{\mu\nu}(x) = L_{\mu}(x)L_{\nu}(x+\mu)L_{\mu}(x+\nu)L_{\nu}(x)$$
(19)

Let us now return to the question of invariance under the local shift transformation

$$\chi(x) \to \alpha_{\lambda}(x+\lambda)\xi_{\lambda}(x)\chi(x+\lambda)$$
(20)

where we have modified eqn. 15 slightly to include change in the argument of $\alpha_{\lambda}(x)$. In this case the transformed action is

$$S' = \sum_{x,\mu} \eta_{\mu}(x)\overline{\chi}(x) \Big[L_{\mu}(x)\alpha_{\lambda}(x)\alpha_{\lambda}(x+\mu)\chi(x+\mu) - L_{\mu}(x-\mu)\alpha_{\lambda}(x-\mu)\alpha_{\lambda}(x)\chi(x-\mu) \Big]$$
(21)

Consider first the case $\lambda \neq \mu$. To ensure invariance of this term requires the addition of new degrees of freedom. Specifically, we can introduce a plaquette (or 2-form) Z_2 gauge field $P_{\mu\lambda}(x)$. Gauge transformations of this object correspond to the appearance of local Z_2 phases for each boundary link of $P_{\mu\lambda}(x)$.

$$P_{\mu\lambda}(x) \to g_{\mu}(x)g_{\lambda}(x+\mu)g_{\mu}(x+\lambda)g_{\lambda}(x)P_{\mu\lambda}(x)$$
(22)

Using this freedom we perform the gauge transformation $P_{\mu\lambda}(x) \rightarrow \alpha_{\lambda}(x)P_{\mu\lambda}(x)\alpha_{\lambda}(x+\mu)$ at the same time as the shift which can then compensate for the factors of $\alpha_{\lambda}(x)\alpha_{\lambda}(x+\mu)$ appearing in the first term on the right hand side of eqn. 21. A similar gauge transformation of $P_{\mu\lambda}(x-\mu)$ can compensate for the factor $\alpha_{\lambda}(x-\mu)\alpha_{\lambda}(x)$ arising in the second term. For this cancellation to work the staggered action must be generalized so that each fermion link term is dressed by the product of all plaquettes that include that link.

$$S = \sum_{x,\mu} \eta_{\mu}(x) \overline{\chi}(x) \left[\left(\prod_{\lambda \neq \mu} P_{\mu\lambda}(x) \right) L_{\mu}(x) \chi(x+\mu) - \left(\prod_{\lambda \neq \mu} P_{\mu\lambda}(x-\mu) \right) L_{\mu}(x-\mu) \chi(x-\mu) \right]$$
(23)

Now we turn to the case $\lambda = \mu$. In this case the action picks up factors like $\alpha_{\mu}(x)\alpha_{\mu}(x \pm \mu)$ corresponding to a shift of two lattice spacings in the μ -direction. This cannot be removed by the transformation of a plaquette (2-form) gauge field $P_{\mu\lambda}(x)$ but can be removed, as described earlier, by a gauge transformation on the link field $L_{\mu}(x)$.

To summarize the gauged action given by eqn. 23 is invariant under the local shift symmetries

$$\chi(x) \to \alpha_{\lambda}(x+\lambda)\xi_{\lambda}(x)\chi(x+\lambda)$$

$$P_{\mu\lambda}(x) \to \delta_{\mu\lambda}P_{\mu\lambda}(x+\lambda) + \alpha_{\lambda}(x+\lambda)P_{\mu\lambda}(x+\lambda)\alpha_{\lambda}(x+\mu+\lambda)\left(1-\delta_{\mu\lambda}\right)$$

$$L_{\lambda}(x) \to \left(1-\delta_{\mu\lambda}\right)L_{\lambda}(x+\lambda) + \alpha_{\lambda}(x+\lambda)L_{\mu}(x+\lambda)\alpha_{\lambda}(x+\mu+\lambda)\delta_{\mu\lambda}$$
(24)

One can add a gauge invariant kinetic term for $P_{\mu\lambda}(x)$ by taking the product of the plaquette fields over all faces of a cube in the lattice and subsequently taking a sum over all cubes. This generalizes the familiar Wilson plaquette operator used for gauge links

$$S_{p} = -\beta_{P} \sum_{x,\mu < \nu < \lambda} F_{\mu\nu\lambda}(x)$$
(25)

where $F_{\mu\nu\lambda}(x)$ is a product over all faces of a cube with corner at x and with basis vectors along the μ , ν and λ directions

$$F_{\mu\nu\lambda}(x) = P_{\mu\nu}(x)P_{\mu\lambda}(x)P_{\nu\lambda}(x)P_{\nu\lambda}(x+\mu)P_{\mu\lambda}(x+\nu)P_{\mu\nu}(x+\lambda)$$
(26)

Taking $\beta_P, \beta_L \to \infty$ forces all the plaquette and link fields to unity up to gauge transformations and the shifts become global symmetries again.

4.2 Higher shifts

If you want to gauge the double shifts one employs the transformations

$$\chi(x) \to \alpha_{\rho\lambda}(x+\rho+\lambda)\xi_{\rho}(x)\xi_{\lambda}(x+\rho)\chi(x+\rho+\lambda)$$
(27)

where $\alpha_{\rho\lambda}(x)$ are Z_2 valued parameters (and $\rho \neq \lambda$). In this case each term in the fermion action changes by a factor of $\alpha_{\rho\lambda}(x)\alpha_{\rho\lambda}(x\pm\mu)$. For each hopping term in direction μ arising in the fermion action there are two cases to consider corresponding to the situation when neither of the two indices ρ and λ equals μ or the case where either ρ or λ equals μ .

Let's first consider the case when $\rho \neq \mu$ and $\lambda \neq \mu$. In this case one can cancel off the factor arising in the transformed fermion kinetic term using a gauge transformation of a cubical (3-form) Z_2 gauge field $C_{\mu\rho\lambda}(x)$ that transforms by local Z_2 factors associated with each $\rho\lambda$ -plaquette face.

$$C_{\mu\rho\lambda}(x) \to \alpha_{\rho\lambda}(x)C_{\mu\rho\lambda}(x)\alpha_{\rho\lambda}(x+\mu)$$
 (28)

In four dimensions there are precisely three such cubes per link. Again a kinetic operator for $C_{\mu\rho\lambda}(x)$ would consist of a product over all eight cubical faces of a hypercube.

$$S_{H} = -\beta_{C} \sum_{x,\mu < \nu < \rho < \lambda} F_{\mu\nu\rho\lambda}(x)$$
⁽²⁹⁾

where

$$F_{\mu\nu\rho\lambda}(x) = C_{\mu\nu\rho}(x)C_{\mu\nu\lambda}(x)C_{\mu\rho\lambda}(x)C_{\nu\rho\lambda}(x)C_{\mu\nu\rho}(x+\lambda)C_{\mu\nu\lambda}(x+\rho)C_{\mu\rho\lambda}(x+\nu)C_{\nu\rho\lambda}(x+\mu)$$
(30)

In the second case one of the directions ρ or λ equals the hopping term index μ . Taking $\rho = \mu$ the kinetic term acquires the factor $\alpha_{\mu\lambda}(x)\alpha_{\mu\lambda}(x \pm \mu)$. There are again three of these terms per link μ . And to retain invariance under such a double shift requires the associated plaquette (2-form) field to transform

$$P_{\mu\lambda}(x) \to \alpha_{\mu\lambda}(x)P_{\mu\lambda}(x)\alpha_{\mu\lambda}(x+\mu)$$
 (31)

The final fermion kinetic term invariant under both single and double shifts now reads

$$S = \sum_{x,\mu} \eta_{\mu}(x) \overline{\chi}(x) \left[\left(\prod_{\lambda,\rho\neq\mu} C_{\mu\rho\lambda}(x) \right) \left(\prod_{\lambda\neq\mu} P_{\mu\lambda}(x) \right) L_{\mu}(x) \chi(x+\mu) - \left(\prod_{\lambda,\rho\neq\mu} C_{\mu\rho\lambda}(x-\mu) \right) \left(\prod_{\lambda\neq\mu} P_{\mu\lambda}(x-\mu) \right) L_{\mu}(x-\mu) \chi(x-\mu) \right]$$
(32)

The corresponding local symmetries are

$$\chi(x) \to a_{\rho\lambda}(x+\rho+\lambda)\xi_{\rho}(x)\xi_{\lambda}(x+\rho)\chi(x+\rho+\lambda)$$

$$C_{\mu\rho\lambda}(x) \to \left(\delta_{\mu\rho}+\delta_{\mu\lambda}\right)C_{\mu\rho\lambda}(x+\rho+\lambda)$$

$$+ a_{\rho\lambda}(x+\rho+\lambda)C_{\mu\rho\lambda}(x+\rho+\lambda)a_{\rho\lambda}(x+\rho+\lambda+\mu)\left(1-\delta_{\mu\rho}-\delta_{\mu\lambda}\right) \qquad (33)$$

$$P_{\rho\lambda}(x) \to \left(1-\delta_{\mu\lambda}-\delta_{\mu\rho}\right)P_{\rho\lambda}(x+\rho+\lambda)$$

$$+ a_{\rho\lambda}(x+\rho+\lambda)P_{\rho\lambda}(x+\rho+\lambda)a_{\rho\lambda}(x+\rho+\lambda+\mu)\left(\delta_{\mu\lambda}+\delta_{\mu\rho}\right) \qquad (34)$$

Clearly this process can be continued to higher shift symmetries using corresponding Z_2 valued lattice p-form gauge fields although clearly there are geometrical obstructions to gauging p-form shifts for p > (D-1) in a D-dimensional lattice. This will be discussed in more detail in the next section.

We should add a caveat at this point. Our strategy for gauging the shift symmetries is to allow the Z_2 -valued α parameters to depend on position. However, we do *not* allow the *type* of shift i.e single, double etc to depend on position. But gauging a discrete group like Γ_4 would usually require such a strategy. Thus our proposal is in some sense only a partial gauging of the discrete symmetry. As we discuss in the next section it will nevertheless yield new and interesting constraints on IR physics.

4.3 Obstructions to gauging - lattice 't Hooft anomalies

Just as the existence of a discrete subgroup of global shift symmetries is sufficient to ensure restoration of the full global flavor symmetry in the continuum limit one hopes that gauging these discrete shift symmetries will lead to a continuum theory with a gauged flavor symmetry.

However, the procedure that we have described using higher form Z_2 valued gauge fields has limitations. Specifically it cannot be used to gauge the γ^5 related shift

$$x \to x + \mu + \nu + \rho + \lambda \quad \mu \neq \nu \neq \rho \neq \lambda \tag{35}$$

since there is no way to write down a 5-cube (5-form) gauge field in four dimensions. And the triple shifts $x \to x + \mu + \lambda + \rho$ (corresponding to the $\gamma^5 \gamma^{\mu}$ generators in the full flavor

group) can be only be subject to a background 4-cube (4-form) gauge field since there is no lattice structure available to carry a kinetic term for this field. Thus the full $\Gamma_4^V \times \Gamma_4^A$ lattice shift symmetry *cannot* be gauged - there is a lattice 't Hooft anomaly. This is consistent with the existence of a continuum 't Hooft anomaly for the $SU(4)_L \times SU(4)_R$ axial-flavor symmetry. Indeed, the existence of a lattice 't Hooft anomaly for Γ_4 strongly suggests that the theory must exhibit a symmetry breaking transition in the continuum limit.

Notice there appears to be no obstruction to gauging a subgroup of the lattice shift symmetries corresponding to single and double shifts using a combination of 0,1 and 2-form Z_2 gauge fields. This would naively correspond to gauging an Sp(4) or SO(4) subgroup of the continuum flavor group. We will examine this possibility in the case of reduced staggered fermions in the next section.

It is interesting to ask what happens to these higher form gauge symmetries in the continuum limit. Since the lattice shifts correspond to discrete flavor transformations the higher form gauge fields must reduce to conventional gauge fields associated with 0-form symmetries in the continuum limit with some of the additional spacetime indices being realized as internal flavor indices. Specifically we conjecture that

$$P_{\mu\lambda}(x) \xrightarrow{a \to 0} P_{\mu}^{\lambda}(x)$$

$$C_{\mu\rho\lambda}(x) \xrightarrow{a \to 0} C_{\mu}^{\rho\lambda}(x)$$
(36)

where the superscript indices are now to be interpreted as operating on the internal flavor space. The non-Abelian character of the continuum flavor symmetry group resides in the properties of the phases $\xi_{\mu}(x)$. As shown in eqn. 5 these phases obey an algebra similar to that of the (Euclidean) gamma matrices and, when these phases are compounded, they mimic the properties of the Γ_4 subgroup of SU(4). We have already seen this in the case of compounding two double shifts and it is easy to see that a similar property holds for a combination of double and single shifts (see appendix)

$$\xi_{\mu}(x+\nu+\lambda)\xi_{\nu}(x+\lambda)\xi_{\lambda}(x)-\xi_{\lambda}(x+\mu+\nu)\xi_{\mu}(x+\nu)\xi_{\nu}(x)=2\left(\delta_{\nu\lambda}\xi_{\mu}(x)-\delta_{\mu\lambda}\xi_{\nu}(x)\right)$$
(37)

which corresponds to a piece of the SU(4) algebra using a Clifford basis for the generators

$$[\gamma^{\mu}\gamma^{\nu},\gamma^{\lambda}] = 2(\delta_{\nu\lambda}\gamma^{\mu} - \delta_{\mu\lambda}\gamma^{\nu})$$
(38)

Notice again that compounding lattice shifts yields another shift in Γ_4 up to a simple translation by a multiple of two lattice spacings – the lattice shift symmetries generate a continuum symmetry group which is a product of both an internal flavor symmetry and the translation group.

In addition, it is necessary to assume some form of universality holds which ensures that the domain of the α variables enhances to the real line in the continuum limit corresponding to the restoration of the full *SU*(4) flavor symmetry. This is certainly expected to be the case in the situation where both shifts and continuum flavor symmetry are both global.

5 Gauging shifts for reduced staggered fermions

Using the site parity operator $\epsilon(x) = (-1)^{\sum_i x_i}$ we can introduce projectors of the form

$$P_{\pm} = \frac{1}{2} \left(1 \pm \epsilon(x) \right) \tag{39}$$

and show that the *massless* staggered fermion action decomposes into two independent parts. Retaining just one of these leads to an action for a *reduced* staggered fermion of the form

$$S = \sum_{x,\mu} \eta_{\mu}(x) \overline{\chi}_{+}(x) [\chi_{-}(x+\mu) - \chi_{-}(x-\mu)]$$
(40)

where $\chi_+ = P_+ \chi$ etc [21,25]. Equivalently we can relabel $\overline{\chi}_+$ as just χ_+ and write

$$S = \sum_{x,\mu} \eta_{\mu}(x)\chi(x)[\chi(x+\mu) - \chi(x-\mu)]$$
(41)

In the continuum matrix theory the projected fields are $\frac{1}{2}(1 \pm \gamma_5 \Psi \gamma_5)$ and are equivalent to reduced Kähler-Dirac fermions [10, 21, 26]. The continuum flavor symmetry is then just $SO(4) \times U(1)$ with the corresponding discrete subgroup of flavor symmetries being

$$G = \{\pm I, \pm i\gamma^{\mu}\gamma^{\nu}, \pm\gamma^5\}$$
(42)

As discussed earlier the maximal symmetry we can gauge is SO(4) and corresponds to gauging the double shifts.

$$\chi(x) \rightarrow i\epsilon(x)\alpha_{\lambda\rho}(x+\lambda+\rho)\xi_{\lambda}(x)\xi_{\rho}(x+\lambda)\chi(x+\lambda+\rho)$$

These shift symmetries correspond to a sixteen element subgroup of SO(4) whose presence guarantees restoration of the global flavor SO(4) symmetry in the continuum limit. These transformations can be gauged by introducing both a link field $L_{\mu}(x)$ and a cube-based Z_2 gauge field as discussed earlier. One again conjectures that this gauged subgroup enhances to the full $SO(4) = SU(2) \times SU(2)$ group in the continuum limit. Notice that the absence of any obstruction to gauging the double shifts in the lattice theory is consistent with the absence of any continuum 't Hooft anomaly for the continuum SO(4) symmetry.

6 Summary

In this paper we have described a proposal to (partially) gauge the shift symmetries of staggered fermions. These shift symmetries correspond to translations within the unit hypercube of a lattice and can be related to a discrete subgroup of the continuum flavor symmetry of four degenerate Dirac fermions ². They give an explicit example of a lattice theory where elementary lattice translations give rise to global internal symmetries in the continuum limit and where lattice 't Hooft anomalies play an important role in understanding the possible phases of the theory [27, 27, 28]. To understand this result it is important to recognize that staggered fermions should be thought of as discrete approximations to Kähler-Dirac fermions whose *twisted* symmetry group corresponds to a diagonal subgroup of Lorentz and flavor symmetries. The shift symmetries then correspond to a discrete subgroup of this twisted rotation group and naturally connect spacetime symmetries with internal symmetries.

The gauging procedure necessitates the introduction of Z_2 -valued higher form gauge fields. However, geometrical obstructions prevent us from gauging the full set of shift symmetries for staggered fermions thereby revealing the presence of a lattice 't Hooft anomaly. This anomaly presumably reflects the existence of the usual continuum 't Hooft anomaly for a theory with an $SU_L(4) \times SU_R(4)$ axial-flavor symmetry.

However, this obstruction is not present in the case of reduced staggered fermions – an observation that is consistent with the absence of anomalies for a theory with just SO(4) flavor

²In this paper we focus exclusively on four (Euclidean) dimensions but clearly the approach we use can be easily generalized to any number of dimensions.

symmetry. This result supports the conjecture that by gauging the double shift symmetry of reduced staggered fermions one can construct a model with emergent SO(4) gauge symmetry.

It was argued in [5, 10] that a continuum theory of reduced Kähler-Dirac fermions is actually chiral and so a successful gauging of the shift symmetries of the corresponding reduced staggered fermion would generate a lattice theory whose continuum limit would describe a particular chiral gauge theory ³. However, this naive expectation will not hold if theory develops couplings between fields on adjacent lattice sites corresponding to mass terms. Such terms are prohibited by the shift symmetry provided it remains a good symmetry under quantization. Our analysis suggests that this is indeed the case.

Acknowledgements

This work was supported by the US Department of Energy (DOE), Office of Science, Office of High Energy Physics, under Award Number DE-SC0009998. SMC would like to acknowledge useful talks with Srimoyee Sen, Shailesh Chandrasekharan, Aleksey Cherman and Yi-Zhuang You.

A Identities

Using

$$\eta_{\mu}(x) = (-1)^{\sum_{i=1}^{\mu-1} x_i}$$
(43)

and

$$\xi_{\lambda}(x) = (-1)^{\sum_{i=\lambda+1}^{4} x_i}, \qquad (44)$$

we see that for $\mu - 1 \ge \lambda$, $\eta_{\mu}(\lambda) = \xi_{\lambda}(\mu) = -1$. And for $\mu - 1 < \lambda$, $\eta_{\mu}(\lambda) = \xi_{\lambda}(\mu) = 1$. This gives the identity

$$\eta_{\mu}(\lambda)\xi_{\lambda}(\mu) = 1. \tag{45}$$

Notice that eqn. 44 implies $\xi_{\mu}(x) = \xi_{\mu}(x + \mu)$, hence $\xi_{\mu}(x)\xi_{\mu}(x + \mu) = 1$. For $\mu \neq \nu$,

$$\xi_{\mu}(x)\xi_{\nu}(x+\mu) + \xi_{\nu}(x)\xi_{\mu}(x+\nu) = (-1)^{\sum_{i=\mu+1}^{4} x_{i}} (-1)^{\sum_{i=\nu+1}^{4} (x+\mu)_{i}} + (\mu \leftrightarrow \nu).$$
(46)

From the above expression it is evident that the two terms differ by a factor of -1 for the two possible cases, $\mu \ge \nu + 1$ and $\nu \ge \mu + 1$. Hence we obtain the identity

$$\xi_{\mu}(x)\xi_{\nu}(x+\mu) + \xi_{\nu}(x)\xi_{\mu}(x+\nu) = 2\delta_{\mu\nu}.$$
(47)

Using eqn. 44, we see that the two terms on the lefthand side of eqn. 37 differ by a factor of $(-1)^{\sum_{i=\mu+1}^{4}\lambda} \times (-1)^{\sum_{i=\nu+1}^{4}\lambda} \times (-1)^{\sum_{i=\lambda+1}^{4}\nu+\mu}$. We restrict to the case $\mu \neq \nu$ to avoid a double shift along the same direction. For $\mu \neq \nu \neq \lambda$, this factor is +1 for all possible orderings, hence the lefthand side of this expression vanishes. For $\mu = \lambda$, this factor simplifies to -1 for the two possible cases, $\nu > \lambda$ or $\nu < \lambda$. Let us first consider the case $\nu > \lambda$, then the first term of eqn. 37 simplifies to

$$\xi_{\mu}(x+\nu+\lambda)\xi_{\nu}(x+\lambda)\xi_{\lambda}(x) = \xi_{\lambda}(x+\nu)\xi_{\nu}(x)\xi_{\lambda}(x) = (-1)^{\sum_{i=\lambda+1}^{4}\nu}\xi_{\nu}(x) = -\xi_{\nu}(x)$$

The second term of Eq. 37 then differs from the first by a factor of -1 as we just argued and hence the equality holds. Repeating the same for $\nu < \lambda$ leads to the same result. For $\nu = \lambda$ the same analysis yields $+2\xi_{\mu}(x)$ on the lefthand side. This proves eqn. 37.

³In fact to cancel off the Z_4 't Hooft anomaly described earlier, the model must contain four reduced fields and then corresponds to the Pati-Salam GUT.

References

- P. Becher and H. Joos, *The Dirac-Kahler Equation and Fermions on the Lattice*, Z. Phys. C 15, 343 (1982), doi:10.1007/BF01614426.
- J. M. Rabin, Homology Theory of Lattice Fermion Doubling, Nucl. Phys. B 201, 315 (1982), doi:10.1016/0550-3213(82)90434-5.
- [3] T. Banks, Y. Dothan and D. Horn, *Geometric Fermions*, Phys. Lett. B117, 413 (1982), doi:10.1016/0370-2693(82)90571-8.
- [4] N. Butt, S. Catterall, A. Pradhan and G. C. Toga, Anomalies and symmetric mass generation for Kähler-Dirac fermions, Phys. Rev. D 104(9), 094504 (2021), doi:10.1103/PhysRevD.104.094504, 2101.01026.
- [5] S. Catterall, 't Hooft anomalies for staggered fermions, Phys. Rev. D 107(1), 014501 (2023), doi:10.1103/PhysRevD.107.014501, 2209.03828.
- [6] E. Kahler, Rend. Math 3-4 **21**, 425 (1962).
- [7] H. B. Nielsen and M. Ninomiya, No Go Theorem for Regularizing Chiral Fermions, Phys. Lett. B 105, 219 (1981), doi:10.1016/0370-2693(81)91026-1.
- [8] S. Catterall, J. Laiho and J. Unmuth-Yockey, *Topological fermion condensates from anomalies*, JHEP 10, 013 (2018), doi:10.1007/JHEP10(2018)013, 1806.07845.
- [9] S. Catterall, J. Laiho and J. Unmuth-Yockey, Kähler-Dirac fermions on Euclidean dynamical triangulations, Phys. Rev. D 98(11), 114503 (2018), doi:10.1103/PhysRevD.98.114503, 1810.10626.
- [10] S. Catterall, Lattice Regularization of Reduced Kähler-Dirac Fermions and Connections to Chiral Fermions, SciPost Phys. 16, 108 (2024), doi:10.21468/SciPostPhys.16.4.108, 2311.02487.
- [11] V. Ayyar and S. Chandrasekharan, *Massive fermions without fermion bilinear condensates*, Phys. Rev. D 91(6), 065035 (2015), doi:10.1103/PhysRevD.91.065035, 1410.6474.
- [12] V. Ayyar and S. Chandrasekharan, Origin of fermion masses without spontaneous symmetry breaking, Phys. Rev. D 93(8), 081701 (2016), doi:10.1103/PhysRevD.93.081701, 1511.09071.
- [13] V. Ayyar and S. Chandrasekharan, *Fermion masses through four-fermion condensates*, JHEP 10, 058 (2016), doi:10.1007/JHEP10(2016)058, 1606.06312.
- [14] S. Catterall, Fermion mass without symmetry breaking, JHEP 01, 121 (2016), doi:10.1007/JHEP01(2016)121, 1510.04153.
- [15] S. S. Razamat and D. Tong, *Gapped Chiral Fermions*, Phys. Rev. X 11(1), 011063 (2021), doi:10.1103/PhysRevX.11.011063, 2009.05037.
- [16] L. Fidkowski and A. Kitaev, The effects of interactions on the topological classification of free fermion systems, Phys. Rev. B 81, 134509 (2010), doi:10.1103/PhysRevB.81.134509, 0904.2197.
- [17] T. Morimoto, A. Furusaki and C. Mudry, Breakdown of the topological classification Z for gapped phases of noninteracting fermions by quartic interactions, Phys. Rev. B92, 125104 (2015), doi:10.1103/PhysRevB.92.125104, 1505.06341.

- [18] Y.-Z. You and C. Xu, Interacting Topological Insulator and Emergent Grand Unified Theory, Phys. Rev. B 91(12), 125147 (2015), doi:10.1103/PhysRevB.91.125147, 1412.4784.
- [19] J. Wang and X.-G. Wen, Nonperturbative definition of the standard model, Phys. Rev. Res.
 2(2), 023356 (2020), doi:10.1103/PhysRevResearch.2.023356, 1809.11171.
- [20] S. Catterall, Chiral lattice fermions from staggered fields, Phys. Rev. D 104(1), 014503 (2021), doi:10.1103/PhysRevD.104.014503, 2010.02290.
- [21] C. van den Doel and J. Smit, Dynamical Symmetry Breaking in Two Flavor SU(N) and SO(N) Lattice Gauge Theories, Nucl. Phys. B228, 122 (1983), doi:10.1016/0550-3213(83)90401-7.
- [22] M. F. L. Golterman and J. Smit, Selfenergy and Flavor Interpretation of Staggered Fermions, Nucl. Phys. B245, 61 (1984), doi:10.1016/0550-3213(84)90424-3.
- [23] M. F. L. Golterman and J. Smit, *Lattice Baryons With Staggered Fermions*, Nucl. Phys. B 255, 328 (1985), doi:10.1016/0550-3213(85)90138-5.
- [24] G. W. Kilcup and S. R. Sharpe, A Tool Kit for Staggered Fermions, Nucl. Phys. B 283, 493 (1987), doi:10.1016/0550-3213(87)90285-9.
- [25] W. Bock, J. Smit and J. C. Vink, Fermion Higgs model with reduced staggered fermions, Phys. Lett. B 291, 297 (1992), doi:10.1016/0370-2693(92)91049-F, hep-lat/9206008.
- [26] W. Bock, J. Smit and J. C. Vink, Staggered fermions for chiral gauge theories: Test on a two-dimensional axial vector model, Nucl. Phys. B 414, 73 (1994), doi:10.1016/0550-3213(94)90422-7, hep-lat/9306012.
- [27] N. Seiberg and S.-H. Shao, Majorana chain and Ising model (non-invertible) translations, anomalies, and emanant symmetries, SciPost Phys. 16, 064 (2024), doi:10.21468/SciPostPhys.16.3.064, 2307.02534.
- [28] M. Cheng and N. Seiberg, Lieb-Schultz-Mattis, Luttinger, and 't Hooft anomaly matching in lattice systems, SciPost Phys. 15(2), 051 (2023), doi:10.21468/SciPostPhys.15.2.051, 2211.12543.