arXiv:2405.06998v1 [math.DG] 11 May 2024

HESSIANIZABILITY OF SURFACE METRICS

ROBERT L. BRYANT

Dedicated to the memory of Joseph A. Wolf, whose generous advice, inspiring works, and seemingly inexhaustible knowledge of geometry has been a boon to me and so many others.

ABSTRACT. A symmetric quadratic form g on a surface M is said to be *locally Hessianizable* if each $p \in M$ has an open neighborhood Uon which there exists a local coordinate chart $(x^1, x^2) : U \to \mathbb{R}^2$ and a function $f: U \to \mathbb{R}$ such that, on U, we have

$$g = \frac{\partial^2 f}{\partial x^i \partial x^j} \, \mathrm{d} x^i \circ \mathrm{d} x^j.$$

In this article, I show that, when g is nondegenerate and smooth, it is always smoothly locally Hessianizable.

Contents

1. Introduction	1
2. The structure equations	3
3. An exterior differential system for Hessianizability	3
3.1. A prolongation	4
4. Involutivity	5
5. Hyperbolicity and smooth Hessianizability	6
5.1. The characteristic variety	6
5.2. Hyperbolicity	7
References	9

1. INTRODUCTION

A symmetric quadratic differential form g on an n-manifold M^n is said to be of *Hessian type* if there exist (n+1) functions x^1, \ldots, x^n, f on M such

Date: January 30, 2024.

²⁰²⁰ Mathematics Subject Classification. 53B05, 58A15.

Key words and phrases. Hessian metrics, Cartan-Kähler theory.

Thanks to Duke University for its support via a research grant, to the NSF for its support via DMS-0103884 and DMS-1359583 (originally DMS-1105868), and to Columbia University for its support via a Samuel Eilenberg Visiting Professorship during Spring 2004, during which this work was begun (and mostly completed).

that $dx^1 \wedge \cdots \wedge dx^n \neq 0$ and such that¹

$$g = \frac{\partial^2 f}{\partial x^i \partial x^j} \, \mathrm{d} x^i \circ \mathrm{d} x^j \,.$$

Note that the independence of the differentials dx^i is needed in order to define the 'partial derivatives' in the formula. One says that g is *locally of Hessian type* if each point of M has an open neighborhood $U \subset M$ on which there exists a coordinate chart $x: U \to \mathbb{R}^n$ and a function $f \in C^{\infty}(U)$ such that the above formula holds on U. The notion of Hessian type has been considered in many places in the literature (cf. [4, 5, 6] and the papers cited therein) and is connected with affine and Kähler geometry and many other areas.

Since symmetric quadratic differential forms g in dimension n depend on $\frac{1}{2}n(n+1)$ functions of n variables and the data of a Hessian representation depends only on (n+1) functions of n variables, it is clear that, when n > 2, not every g is locally of Hessian type. As n increases, the condition of being locally of Hessian type becomes more and more restrictive. When g is nondegenerate, so that (M^2, g) is a pseudo-Riemannian manifold, being of Hessian type implies algebraic conditions on the Riemann curvature tensor once n is sufficiently large [6, 1].

Meanwhile, when n = 2, finding Hessianizing data (x^1, x^2, f) for a given quadratic form g is a determined PDE problem, being three equations for three unknowns, so one might expect formal solvability. However, it is not necessarily true that the PDE system involved is nondegenerate, so even formal solvability is not guaranteed.

For example, the smooth (in fact, real-analytic) quadratic form $g = c_{ijk}y^i dy^j \circ dy^k$, where $c_{ijk} = c_{ikj}$ are constants, is not Hessianizable on a neighborhood of y = 0 unless $c_{ijk} = c_{jik}$.

However, when $g \in S^2(T^*M)$ is real-analytic and nondegenerate at $p \in M^2$, it is straightforward to show, using the Cartan-Kähler Theorem, that g is Hessianizable on an open neighborhood of p. See Theorem 1 below, which is the result to which I alluded in [3]. (The reader might also want to compare the argument in [1], which was arrived at independently.)

It is natural to ask whether this Hessianizability result continues to hold in the smooth setting. It turns out that the PDE involved is never elliptic, so one never gets existence and/or regularity by applying results from elliptic theory. In fact, the complex characteristic variety of the PDE along regular initial conditions consists of 3 points, so at least one of the characteristics has to be real.

Now, it turns out that, when g is nondegenerate, one can always find regular initial conditions for the PDE with 3 real, distinct characteristics, i.e., initial conditions for which the PDE problem is hyperbolic. Thus, one

¹Note the use of the Einstein summation convention, which will be employed throughout this article. Also, note the standard use of $\alpha \circ \beta$ for $\frac{1}{2}(\alpha \otimes \beta + \beta \otimes \alpha)$, i.e., the symmetrized tensor product.

finds that one can invoke results of Yang [7] on involutive hyperbolic systems to prove smooth local Hessianizability of nondegenerate g that are merely smooth. See Theorem 2.

For details on the notation and definitions in the use of the Cartan-Kähler Theorem, which will be used to prove Theorem 1, the reader is referred to [2].

2. The structure equations

Let M^2 be a surface, and let g be a metric on M, i.e., a nondegenerate quadratic form on M. Thus, in a local coordinate chart $y = (y^i) : U \to \mathbb{R}^2$, one has $g = g_{ij}(y) dy^i dy^j$, where $g_{11}g_{22} - g_{12}^2$ is nonvanishing. (One does not need to assume that g be definite.)

Let $\pi : F \to M$ denote the coframe bundle of M, whose typical element $u \in F$ is an isomorphism $u : T_{\pi(u)}M \to \mathbb{R}^2$, with its right action by $A \in \operatorname{GL}(2,\mathbb{R}) = \operatorname{Aut}(\mathbb{R}^2)$ defined to be $R_A(u) = u \cdot A = A^{-1} \circ u$. There is an \mathbb{R}^2 -valued 1-form $\omega : TF \to \mathbb{R}^2$ defined by $\omega_u(v) = u(\pi'(v))$ for $v \in T_uF$. Its components $\omega = (\omega^i)$ are the canonical ('soldering') forms ω^i (where $1 \leq i \leq 2$). Let $G_{ij} = G_{ji}$ be the functions on F such that $\pi^*g = G_{ij}\omega^i \circ \omega^j$ and, as usual, let $G^{ij} = G^{ji}$ satisfy $G^{ik}G_{kj} = \delta^i_j$.

Let ω_j^i be the Levi-Civita connection forms of g on F. These forms, together with the ω^i , define a coframing of F, and they are uniquely specified by the requirements that they satisfy *compatibility* with g,

(2.1)
$$\mathrm{d}G_{ij} = G_{ik}\,\omega_j^k + G_{kj}\,\omega_i^k\,,$$

and the first structure equation,

(2.2)
$$\mathrm{d}\omega^i = -\omega^i_i \wedge \omega^j.$$

Furthermore, they satisfy the second structure equation,

(2.3)
$$d\omega_j^i = -\omega_k^i \wedge \omega_j^k + G_{jl} K \,\omega^i \wedge \omega^l$$

where the function K is the Gauss curvature of g pulled up to F via the projection $\pi: F \to M$.

It will be convenient to define $\omega_{ij} = G_{ik} \omega_j^k$, so that the equations above become $dG_{ij} = \omega_{ij} + \omega_{ji}$, and to define $\omega_i = G_{ij} \omega^j$, so that $\pi^*g = \omega_i \circ \omega^i$. Note that one has $\omega_i \wedge \omega^i = 0$, since $G_{ij} = G_{ji}$.

3. An exterior differential system for Hessianizability

If $U \subset M$ is an open subset on which there exist coordinates $x = (x^i)$: $U \to \mathbb{R}^2$ and a function f such that, on U, one has

$$g = \frac{\partial^2 f}{\partial x^i \partial x^j} \, \mathrm{d} x^i \circ \mathrm{d} x^j = \mathrm{d} \left(\frac{\partial f}{\partial x^1} \right) \circ \mathrm{d} x^1 + \mathrm{d} \left(\frac{\partial f}{\partial x^2} \right) \circ \mathrm{d} x^2,$$

then the coordinate coframing $\xi = (\mathrm{d}x^i)$ defines a section $\xi : U \to F$ that satisfies $\xi^*(\omega^i) = \mathrm{d}x^i$ and $\xi^*(G_{ij}) = \frac{\partial^2 f}{\partial x^i \partial x^j}$. Consequently, $\xi^*(\omega_i) = \mathrm{d}\left(\frac{\partial f}{\partial x^i}\right)$. In particular, the section $\xi: U \to F$ satisfies

$$\xi^*(\mathrm{d}\omega^i) = \mathrm{d}(\mathrm{d}x^i) = 0 \quad \text{and} \quad \xi^*(\mathrm{d}\omega_i) = \mathrm{d}\left(\mathrm{d}\left(\frac{\partial f}{\partial x^i}\right)\right) = 0$$

while $\xi^*(\omega^1 \wedge \omega^2) = \mathrm{d}x^1 \wedge \mathrm{d}x^2 \neq 0.$

Conversely ², if $S \subset F$ is a surface such that $S^*(d\omega^i) = S^*(d\omega_i) = 0$ while $S^*(\omega^1 \wedge \omega^2) \neq 0$, then each point of S has an open neighborhood $V \subset S$ that is the image of a local section $\xi : U \to F$ for some open set $U \subset M$. When U is simply connected, one has $\xi^*\omega^i = dx^i$ for some independent functions x^i on U, while $\xi^*\omega_i = dp_i$ for some functions p_i on U. Since $\omega_i \wedge \omega^i = 0$, it follows that $d(p_i dx^i) = dp_i \wedge dx^i = 0$, so that there exists a function f on U such that $df = p_i dx^i$, i.e., $p_i = \frac{\partial f}{\partial x^i}$. Thus,

$$g = \xi^*(\omega_i \circ \omega^i) = \mathrm{d}p_i \circ \mathrm{d}x^i = \frac{\partial^2 f}{\partial x^i \partial x^j} \,\mathrm{d}x^i \circ \mathrm{d}x^j,$$

as desired.

Letting \mathcal{I} be the differential ideal on F generated by the four 2-forms $d\omega^i$ and $d\omega_i$ for i = 1, 2, it then follows that the problem of locally writing g in Hessian form is equivalent to finding the integral surfaces of \mathcal{I} on which the 2-form $\omega^1 \wedge \omega^2$ is nonvanishing.

3.1. A prolongation. The ideal with independence condition $(\mathcal{I}, \omega^1 \wedge \omega^2)$ is not quite the right ideal to consider, for it does not contain all of the forms on F that vanish on Hessianizing sections. To see this, note that, by definition, $G_{ij}d\omega^j = -\omega_{ij}\wedge\omega^j$ while, since $dG_{ij} = \omega_{ij} + \omega_{ji}$, one has that

$$\mathrm{d}\omega_i = \mathrm{d}G_{ij} \wedge \omega^j + G_{ij}\mathrm{d}\omega^j = \omega_{ji} \wedge \omega^j.$$

Thus, \mathcal{I} is generated by the four 2-forms $\omega_{ij}\wedge\omega^j$ and $\omega_{ji}\wedge\omega^j$ for i = 1, 2. In particular, \mathcal{I} contains the pair of 2-forms $(\omega_{12} - \omega_{21})\wedge\omega^j$ for j = 1, 2, and, consequently, any integral surface of \mathcal{I} on which $\omega^1\wedge\omega^2 \neq 0$ must also be an integral surface of the 1-form $\theta = \frac{1}{2}(\omega_{12} - \omega_{21})$. Thus, one can add θ and its exterior derivative $d\theta$ to \mathcal{I} to form a larger ideal \mathcal{I}_+ with the property that the integral surfaces of $(\mathcal{I}_+, \omega^1 \wedge \omega^2)$ are the same as the integral surfaces of $(\mathcal{I}, \omega^1 \wedge \omega^2)$. Henceforth, I will work with this larger ideal, which is a partial prolongation of $(\mathcal{I}, \omega^1 \wedge \omega^2)$.

Write $\sigma_{ij} = \frac{1}{2}(\omega_{ij} + \omega_{ji}) \ (= \frac{1}{2} dG_{ij})$, so that $\omega_{ij} \equiv \sigma_{ij} \mod \theta$ and $-G_{ij} d\omega^j \equiv d\omega_i \equiv \sigma_{ij} \wedge \omega^j \mod \theta$.

Note that the forms ω^1 , ω^2 , θ , σ_{11} , σ_{12} , and σ_{22} are everywhere linearly independent and hence define a coframing of F. Using the second structure

4

²Here and throughout this article, when $S \subset M$ is a submanifold and τ is a contravariant tensor field on M (e.g., a differential form or a metric on M), the notation $S^*(\tau)$ denotes the *pullback* of τ to S, which the reader should take care not to confuse with $\tau_{|S}$, i.e., the *restriction* of τ to S.

equations, one finds

$$\mathrm{d}\theta \equiv G^{kl}\sigma_{k1} \wedge \sigma_{l2} + K\,\omega_1 \wedge \omega_2 \mod \theta.$$

Thus, the differential ideal \mathcal{I}_+ is generated algebraically by θ and three 2-forms:

(3.1)
$$\Omega_1 = \sigma_{1j} \wedge \omega^j,$$
$$\Omega_2 = \sigma_{2j} \wedge \omega^j,$$
$$\Theta = G^{kl} \sigma_{k1} \wedge \sigma_{l2} + K \omega_1 \wedge \omega_2$$

It is easy to see that the right action of $\operatorname{GL}(2,\mathbb{R})$ on F preserves the ideal with independence condition $(\mathcal{I}_+, \omega^1 \wedge \omega^2)$. In particular, this right action prolongs to an action of $\operatorname{GL}(2,\mathbb{R})$ on the space $\mathcal{V}_2(\mathcal{I}_+, \omega^1 \wedge \omega^2) \subset \operatorname{Gr}_2(TF)$, which consists of the 2-dimensional integral elements of \mathcal{I}_+ on which $\omega^1 \wedge \omega^2$ is nonvanishing.

4. Involutivity

The pieces are now in place to prove the main result.

Theorem 1. The differential ideal with independence condition $(\mathcal{I}_+, \omega^1 \wedge \omega^2)$ is involutive, with Cartan characters $s_0 = 1$, $s_1 = 3$, and $s_2 = 0$.

If g is real analytic, then the integral surfaces of $(\mathcal{I}_+, \omega^1 \wedge \omega^2)$ depend on three arbitrary functions of one variable. In particular, each point of M has an open neighborhood on which g can be written in Hessian form.

Proof. Because \mathcal{I}_+ is generated in degree 1 by the nonvanishing 1-form θ , all of the 0-dimensional integral elements are both ordinary and regular, and all of its 1-dimensional integral elements are ordinary.

Thus, to prove involutivity, it suffices to show that, at each $u \in F$, there exists a regular 1-dimensional integral element $E_1 \subset T_u F$ for which $\omega^1 \wedge \omega^2$ is nonvanishing on the polar space $H(E_1)$.

To this end, let $v \in T_u F$ be the basis of a 1-dimensional integral element, i.e., $\theta(v) = 0$ while v is nonzero. Then the polar space of $E_1 = \mathbb{R}v$ is

$$H(E_1) = \{ w \in T_u F \mid \theta(w) = \Omega_1(v, w) = \Omega_2(v, w) = \Theta(v, w) = 0 \},\$$

and hence has dimension at least 2. Set $a^i = \omega^i(v)$ and $s_{ij} = \sigma_{ij}(v)$. Then

$$v \, \lrcorner \, \Omega_1 = -a^1 \, \sigma_{11} - a^2 \, \sigma_{12} + s_{11} \, \omega^1 + s_{12} \, \omega^2,$$

(4.1) $v \lrcorner \Omega_2 = -a^1 \sigma_{12} - a^2 \sigma_{22} + s_{12} \omega^1 + s_{22} \omega^2,$

$$v \,\lrcorner\, \Theta = G^{kl}(s_{k1}\sigma_{l2} - s_{l2}\sigma_{k1}) + K(G_{11}G_{22} - G_{12}^{2})\,(a^1\,\omega^2 - a^2\,\omega^1).$$

Thus, $H(E_1)$ has dimension 2 and $\omega^1 \wedge \omega^2$ is nonvanishing on $H(E_1)$ exactly when

(4.2)
$$(a^1)^2 s_1^2 - a^1 a^2 (s_1^1 - s_2^2) - (a^2)^2 s_2^1 \neq 0,$$

where, for brevity, I have written $s_j^i = G^{ik}(u)s_{kj}$. In particular, except when $s_{ij} = \lambda G_{ij}(u)$ for some λ , it will always be possible to choose a^1 and a^2 so

R. BRYANT

that the inequality (4.2) holds. For a $v \in T_u F$ so chosen, the integral element $E_1 = \mathbb{R}v$ is regular and, hence, its 2-dimensional polar space $H(E_1) = E_2$ is an ordinary integral element of $(\mathcal{I}_+, \omega^1 \wedge \omega^2)$. Moreover, the characters of the ordinary flag $(0) \subset E_1 \subset E_2$ are visibly $s_0 = 1$, $s_1 = 3$, and $s_2 = 0$ and $(\mathcal{I}_+, \omega^1 \wedge \omega^2)$ is involutive.

Moreover, when g is real analytic, the Cartan-Kähler Theorem applies, yielding that any analytic curve $\Gamma \subset F$ that satisfies $\Gamma^*\theta = 0$ and $\Gamma^*C \neq 0$, where C is the characteristic cubic form³

(4.3)
$$C = (\omega^1)^2 \circ (G^{2j}\sigma_{j1}) - \omega^1 \circ \omega^2 \circ (G^{1j}\sigma_{j1} - G^{2j}\sigma_{j2}) - (\omega^2)^2 \circ (G^{1j}\sigma_{j2}),$$

lies in a unique (ordinary) analytic integral surface of $(\mathcal{I}_+, \omega^1 \wedge \omega^2)$.

When g (and hence \mathcal{I}_+) is real analytic, it is easy to construct such analytic curves $\Gamma \subset F$: Let $A \subset M$ be any connected, noncompact, analytic, embedded curve, on which there is a parameter $t : A \to (a, b) \subset \mathbb{R}$ and let $F_A = \pi^{-1}(A) \subset F$. Then F_A is a 5-dimensional embedded, analytic submanifold on which $\omega^1 \wedge \omega^2$ vanishes, although ω^1 and ω^2 do not simultaneously vanish. Thus, there exist two functions $a^i : F_A \to \mathbb{R}$ for i = 1, 2that do not simultaneously vanish such that $\omega^i = a^i \pi^*(dt)$ on F_A . Then let $\Gamma \subset F_A$ be any real analytic curve in F_A that projects diffeomorphically onto A and satisfies $\Gamma^*\theta = 0$ while $\Gamma^*\psi \neq 0$ where ψ is the 1-form on F_A defined by

$$\psi = (a^1)^2 G^{2k} \sigma_{k1} - a^1 a^2 (G^{1k} \sigma_{k1} - G^{2k} \sigma_{k2}) - (a^2)^2 G^{1k} \sigma_{k2}.$$

Such a Γ will then be a regular integral curve of \mathcal{I}_+ lying in a unique integral surface $S \subset F$ of $(\mathcal{I}_+, \omega^1 \wedge \omega^2)$. Moreover, an open neighborhood of Γ in Swill project diffeomorphically onto an open neighborhood U of A in M, and the inverse map $\xi : U \to S$ will be a Hessianizing section for g on U.

In particular, for any given point $u \in F$, an integral surface of $(\mathcal{I}_+, \omega^1 \wedge \omega^2)$ exists that passes through u.

Since the Cartan characters of the system are $s_0 = 1$, $s_1 = 3$, and $s_2 = 0$, it follows that the ordinary integral surfaces depend on three functions of one variable [2].

As explained above, this implies that the metric g is locally Hessianizable if it is real analytic.

5. Hyperbolicity and smooth Hessianizability

In order to prove Hessianizability in the smooth category, it will be necessary to take a closer look at the structure of the ideal with independence condition $(\mathcal{I}_+, \omega^1 \wedge \omega^2)$.

5.1. The characteristic variety. Let $\mathcal{V}_2(\mathcal{I}_+, \omega^1 \wedge \omega^2) \subset \operatorname{Gr}_2(TF)$ denote the locus of 2-dimensional integral elements of $(\mathcal{I}_+, \omega^1 \wedge \omega^2)$. A 2-plane $E \subset$

³For use below, I note that $R_A^*(C) = \det(A) C$ for $A \in \operatorname{GL}(2, \mathbb{R})$.

 $T_u F$ belongs to $\mathcal{V}_2(\mathcal{I}_+, \omega^1 \wedge \omega^2)$ if and only if it is defined by 4 Pfaffian equations of the form

(5.1)
$$\theta = \sigma_{ij} - p_{ijk}(E)\,\omega^k = 0,$$

where the numbers $p_{ijk}(E) = p_{jik}(E)$ satisfy the linear relations $p_{ijk}(E) = p_{ikj}(E)$ (which are equivalent to the vanishing of Ω_i on E) and the quadratic relation

(5.2)
$$0 = G^{kl}(u) \left(p_{k11}(E) p_{l22}(E) - p_{k12}(E) p_{l12}(E) \right) + K(u) \left(G_{11}(u) G_{22}(u) - G_{12}(u)^2 \right)$$

(which is equivalent to the vanishing of Θ on E). The quadratic form in the 4 independent quantities $p_{111}(E)$, $p_{112}(E)$, $p_{122}(E)$, and $p_{222}(E)$ that makes up the first term of this relation is easily seen to be nondegenerate and indefinite. Hence, this exhibits $\mathcal{V}_2(\mathcal{I}_+, \omega^1 \wedge \omega^2)$ as a hypersurface in $F \times \mathbb{R}^4$ whose fiber over $u \in F$ is a (nonempty) 3-dimensional hyperquadric that is smooth away from the 2-plane defined by $p_{ijk}(E) = K(u) = 0$.

Meanwhile, the integral element E will be ordinary (i.e., it will contain a regular 1-dimensional integral element) if and only if $E^*(C) \neq 0$, since $\Xi_E \subset \mathbb{P}E^*$, i.e., the characteristic variety of E, consists of the points represented by the linear factors of the cubic form

(5.3)
$$E^*(C) = \epsilon_{jk} G^{kl}(u) p_{lmn}(E) \ \omega^j \circ \omega^m \circ \omega^n.$$

(Here, $\epsilon_{ij} = -\epsilon_{ji}$ and $\epsilon_{12} = 1$.) It follows that E is ordinary unless $p_{ijk}(E) = 0$ for all (i, j, k) which, by the above quadratic relation, can only happen if K(u) = 0.

Remark 1 (Non-ordinary integral surfaces). An integral surface of the ideal $(\mathcal{I}_+, \omega^1 \wedge \omega^2)$ on which C vanishes identically is a so-called 'non-ordinary' integral surface, i.e., it is an integral surface of $(\mathcal{I}_+, \omega^1 \wedge \omega^2)$ that contains no regular integral curves. Since $E^*(C)$ vanishes if and only if $p_{ijk}(E) = 0$, it follows from (5.2) that non-ordinary integral surfaces can only exist over open sets in M^2 on which the curvature K vanishes, and, in this case, they are the integral surfaces of the Frobenius system generated by θ and the σ_{ij} . Thus, when $K \equiv 0$, the coframe bundle F is foliated by the 4-parameter family of leaves of this system.

Conversely, if g is a nondegenerate quadratic form on M with vanishing curvature, then each point $p \in M$ has a neighborhood on which there exist p-centered local coordinates (x^1, x^2) in which $g = c_{ij} dx^i dx^j$, where the $c_{ij} = c_{ji}$ are constants, so that g is in Hessian form with $f = \frac{1}{2}c_{ij}x^ix^j$. Such constant coefficient Hessianizations describe the non-ordinary integral surfaces.

5.2. Hyperbolicity. The (complex) characteristic variety of the ideal \mathcal{I}_+ meets each cotangent space of each ordinary integral surface in the three linear factors of the cubic form C (as defined in (4.3)). Since a nonvanishing cubic in 2 variables always has at least one real linear factor, it follows that

R. BRYANT

the real characteristic variety is never empty, so the system is never elliptic. Therefore, the best behavior one could hope for is that the characteristic variety at each point would consist of three real distinct factors, which would be the hyperbolic case.

Recall that, for a cubic form $c \in S^3(E^*)$, where E is a 2-dimensional vector space over \mathbb{R} , its discriminant $\Delta(c)$ is an element of $S^6(\Lambda^2(E^*))$. When (e^1, e^2) is a basis of E^* and c is expressed in the form

(5.4)
$$c = c_0 (e^1)^3 + 3c_1 (e^1)^2 e^2 + 3c_2 e^1 (e^2)^2 + c_3 (e^2)^3$$

then

(5.5)
$$\Delta(c) = \left(6c_0c_1c_2c_3 + 3c_1^2c_2^2 - 4c_0c_2^3 - 4c_1^3c_3 - c_0^2c_3^2\right)(e^1 \wedge e^2)^{\otimes 6}.$$

Then $\Delta(c) = 0$ if and only if c has a multiple linear factor. Because the scalar field is \mathbb{R} , the inequality $\Delta(c) > 0$ makes sense, and this inequality holds if and only if c has three real, distinct, linear factors.

Let $\mathcal{V}_2^h(\mathcal{I}_+, \omega^1 \wedge \omega^2) \subset \mathcal{V}_2(\mathcal{I}_+, \omega^1 \wedge \omega^2)$ denote the open subset that consists of integral elements E for which the characteristic cubic $E^*(C)$ is nonzero and has three real, distinct, linear factors. By definition, the elements of $\mathcal{V}_2^h(\mathcal{I}_+, \omega^1 \wedge \omega^2)$ are ordinary. I will refer to $\mathcal{V}_2^h(\mathcal{I}_+, \omega^1 \wedge \omega^2)$ as the hyperbolic locus.

Lemma 1. The basepoint projection $\mathcal{V}_2^h(\mathcal{I}_+, \omega^1 \wedge \omega^2) \to M$ is a surjective submersion.

Proof. It suffices to prove that the basepoint projection $\mathcal{V}_2^h(\mathcal{I}_+, \omega^1 \wedge \omega^2) \rightarrow M$ is surjective, since every element of $\mathcal{V}_2^h(\mathcal{I}_+, \omega^1 \wedge \omega^2)$ is a Kähler-regular integral element and the ideal \mathcal{I}_+ is generated in positive degree.

Also, because the right action by $GL(2, \mathbb{R})$ on F is transitive on the fibers of $F \to M$, it suffices to consider an element $u \in F$ at which all of G_{11} , G_{12} and G_{22} are nonzero, so I will assume this hencforth.

Consider the locus $L_u \subset \mathcal{V}_2(\mathcal{I}_+, \omega^1 \wedge \omega^2)$ in the fiber over u that is defined by the two equations

$$G_{22}p_{122} - G_{12}p_{222} = 0$$
 and $G_{12}p_{111} - G_{11}p_{112} = 0$.

Because $G_{11}G_{22} - (G_{12})^2 \neq 0$, there will exist unique functions s_1 and s_2 on L such that

$$p_{111} = G_{11}s_1, \ p_{112} = G_{12}s_1, \ p_{122} = G_{12}s_2, \ p_{222} = G_{22}s_2$$

These values for p_{ijk} will define an integral element if and only if they satisfy (5.2), which turns out to be

(5.6)
$$(G_{11}G_{22}-G_{12}^2)^2 K - G_{12}(G_{11}s_2-G_{12}s_1)(G_{21}s_2-G_{22}s_1) = 0.$$

Because G_{12} and $G_{11}G_{22} - G_{12}^2$ are nonvanishing, this equation defines either a hyperbola or a pair of transverse lines in the s_1s_2 -plane, depending on whether K is zero or not.

8

Meanwhile, the characteristic cubic at points of L_u is (up to a nonzero multiple) given by

$$C = -((G_{11}G_{22} + G_{12}^2)s_1 - 2G_{11}G_{12}s_2)(\omega^1)^2\omega^2 + ((G_{11}G_{22} + G_{12}^2)s_2 - 2G_{22}G_{12}s_1)\omega^1(\omega^2)^2.$$

Obviously C will have three real distinct factors as long as both quantities $(G_{11}G_{22} + G_{12}^2)s_1 - 2G_{11}G_{12}s_2$ and $(G_{11}G_{22} + G_{12}^2)s_2 - 2G_{22}G_{12}s_1$ are nonzero. The vanishing of these quantities define two lines ℓ_1 and ℓ_2 in the s_1s_2 -plane that intersect transversely at $s_1 = s_2 = 0$.

When K is nonzero, the hyperbola (5.6) in the s_1s_2 -plane is not contained in the union of the two lines ℓ_1 and ℓ_2 . Hence there is a choice of (s_1, s_2) that satisfies (5.6) for which C has three real distinct factors.

When K = 0, one checks that the two (distinct) lines in the s_1s_2 -plane defined by $G_{11}s_2-G_{12}s_1 = 0$ and $G_{21}s_2-G_{22}s_1 = 0$ only meet the lines ℓ_1 and ℓ_2 in the origin $s_1 = s_2 = 0$. Hence, again there is a choice of (s_1, s_2) that satisfies (5.6) for which C has three real distinct factors.

Such a choice of (s_1, s_2) defines an element of L_u that lies in $\mathcal{V}_2^h(\mathcal{I}_+, \omega^1 \wedge \omega^2)$, proving the claimed surjectivity.

Remark 2 (Universal hyperbolicity). It is not always true that each $u \in F$ has a 2-dimensional integral element $E \subset T_u F$ for which the characteristic cubic $E^*(C)$ is the product of a linear factor and an irreducible quadratic factor (the other open $\operatorname{GL}(2,\mathbb{R})$ -orbit). For example, if g > 0, then such integral elements exist only when K(u) < 0. In fact, when g > 0 and K > 0, one finds that *all* of the Kähler-ordinary integral elements are hyperbolic.

Theorem 2. Let $\mathcal{I}^{(1)}$ denote the contact ideal on $\mathcal{V}_2^h(\mathcal{I}_+, \omega^1 \wedge \omega^2) \subset \operatorname{Gr}_2(TF)$, endowed with its canonical independence condition ω . Then $(\mathcal{I}^{(1)}, \omega)$ is involutive hyperbolic in the sense of Yang [7].

Consequently, any smooth nondegenerate quadratic form on M^2 is locally Hessianizable.

Proof. Since $(\mathcal{I}^{(1)}, \omega)$ on $\mathcal{V}_2^h(\mathcal{I}_+, \omega^1 \wedge \omega^2)$ is an open subset of the ordinary prolongation of the involutive system $(\mathcal{I}_+, \omega^1 \wedge \omega^2)$, it is involutive, with Cartan characters $s_0 = 4$, $s_1 = 3$, $s_2 = 0$. Moreover, because ordinary prolongation preserves the characteristic variety, it follows that the characteristic variety of each integral element of $(\mathcal{I}^{(1)}, \omega)$ consists of three real, distinct points. Thus, $(\mathcal{I}^{(1)}, \omega)$ is involutive hyperbolic in the sense of Yang [7].

In particular, it follows from Theorem 4.11 in [7] that, when g is a smooth nondegenerate quadratic form on M^2 , each $u \in F$ lies in a smooth, ordinary, *hyperbolic* integral surface of $(\mathcal{I}_+, \omega^1 \wedge \omega^2)$. Consequently, any smooth nondegenerate quadratic form on a surface can locally be written in Hessian form.

References

 S. Amari and J. Armstrong, *Curvature of Hessian manifolds*, Differential Geometry and Its Applications, **33** (2014), 1–12.

R. BRYANT

- [2] R. Bryant, S.-s. Chern, R. Gardner, H. Goldschmidt, and P. Griffiths, *Exterior dif*ferential systems, Springer-Verlag, New York, 1991.
- [3] R. Bryant, Answer to mathoverflow question "When a Riemannian manifold is of Hessian Type", http://mathoverflow.net/questions/122308/.
- [4] J. Duistermaat, On Hessian Riemannian structures, Asian J. Math. 5 (2001), 79–91.
- [5] H. Shima, *The geometry of Hessian structures*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2007.
- [6] B. Totaro, The curvature of a Hessian metric, Int. J. Math. 15 (2004), 369–391.
- [7] D. Yang, Involutive hyperbolic differential systems. Mem. Amer. Math. Soc. 68 (1987), no. 370.

DUKE UNIVERSITY MATHEMATICS DEPARTMENT, P.O. BOX 90320, DURHAM, NC 27708-0320

Email address: bryant@math.duke.edu

URL: http://www.math.duke.edu/~bryant